

Read questions carefully before answering. No outside paper is allowed. Write **set up equation** for a mathematical problem, then put the raw data with **units**, before showing the calculation. Use the reverse side of your answer paper as scratch. Use the periodic table and important constants charts provided. (Total points = $66 + (16 \times 3) = 114$).

Show your calculation with set up and units (when appropriate)

- 1) The initial rate of the reaction $A + B \rightarrow C$ was measured at several different concentrations of the reactants. Following **formal** methods, (a) calculate the rate law for the reaction (6 pts.) and (b) The magnitude of the rate constant (2 pts.). 1) _____

Experiment	Initial Concentrations		Initial Rate ($M s^{-1}$)
	[A] (M)	[B] (M)	
1	0.10	0.10	4.0×10^{-5}
2	0.10	0.20	4.0×10^{-5}
3	0.20	0.10	16.0×10^{-5}

- 2) In the reaction $N_2(g) + 3 H_2(g) \leftrightarrow 2 NH_3(g)$, if the $K_C = 9.60$ at 573 K, then calculate the K_p at this temperature (4 pts.). 2) _____

3) Activation energies of reactions, E_a , are frequently found graphically. The Arrhenius equation: $\ln(k) = (-E_a)/RT + \ln(A)$ is used. Values of k , the rate constant, are measured at various temperatures, then $\ln k$ and $1/T$ are calculated and plotted. In one particular experiment the **slope** of the st. line (obtained by plotting $\ln(k)$ and $1/T$) was found to be -30000 K . Calculate the energy of activation (in calories) of the reaction (6 pts.):

3) _____

4) Equilibrium was established when a mixture of 0.20 mol of NO(g) , 0.10 mol of $\text{H}_2\text{(g)}$, and 0.20 mol of $\text{H}_2\text{O(g)}$ is placed in a 2.0-L vessel at 400 K. The equilibrium reaction is : $2 \text{NO(g)} + 2 \text{H}_2\text{(g)} \rightleftharpoons \text{N}_2\text{(g)} + 2 \text{H}_2\text{O(g)}$. If at equilibrium $[\text{NO}] = 0.062 \text{ M}$, then calculate K_p . (10 pts.)

4) _____

5) Calculate the pOH of a solution if 1.35 moles of HI is in 530.00 mL of water. (6 pts.)

5) _____

6) Ammonia is a weak base with $pK_b = 4.74$ at 25°C . Calculate the pH of a 0.2 M ammonia solution in water at that temperature (8 pts.). 6) _____

7) Calculate the pH of a buffer solution that contains 0.820 grams of sodium acetate and 0.01 moles of acetic acid in 100 ml of water. The K_a of acetic acid is 1.77×10^{-5} (8 pts). 7) _____



8) K_{sp} of Ag_2CrO_4 in water at 20°C is 1.9×10^{-12} M. Calculate its solubility in gram per 0.1 L of solution (8 pts.). 8) _____

- 9) Calculate the molar solubility of CaF_2 at 25°C in a solution that is 0.010 M in $\text{Ca}(\text{NO}_3)_2$. 9) _____
K_{sp} for $\text{CaF}_2 = 3.9 \times 10^{-11}$. Show your calculation with ICE chart. (8 pts.)

MULTIPLE CHOICE. On your scantron, start answering from number as the question number. Select the one alternative that best completes the statement or answers the question (3 pts each).


- 10) The rate of a reaction depends on _____. 10) _____
A) collision energy
B) collision orientation
C) collision frequency
D) all of the above
E) none of the above
- 11) A catalyst can increase the rate of a reaction _____. 11) _____
A) by providing an alternative pathway with a lower activation energy
B) by changing the value of the frequency factor (A)
C) by increasing the overall activation energy (E_a) of the reaction
D) by lowering the activation energy of the reverse reaction
E) All of these are ways that a catalyst might act to increase the rate of reaction.
- 12) How does the reaction quotient of a reaction (Q) differ from the equilibrium constant (K_{eq}) of the same reaction? 12) _____
A) K does not depend on the concentrations or partial pressures of reaction components.
B) Q is the same as K_{eq} when a reaction is at equilibrium.
C) Q does not depend on the concentrations or partial pressures of reaction components.
D) K_{eq} does not change with temperature, whereas Q is temperature dependent.
E) Q does not change with temperature.

13) The equilibrium constant for reaction 1 is K. The equilibrium constant for reaction 2 is _____.



- A) $1/2K$ B) K^2 C) $-K^2$ D) $2K$ E) $1/K^2$

14) The reaction below is exothermic:



Le Chatelier's Principle predicts that _____ will result in an increase in the number of moles of $\text{SO}_3(\text{g})$ in the reaction container.

- A) removing some oxygen
 B) increasing the pressure
 C) increasing the volume of the container
 D) decreasing the pressure
 E) increasing the temperature

15) The equilibrium reaction $\text{Co}(\text{H}_2\text{O})_6^{2+}(\text{aq})$ (Pink) + $4 \text{Cl}^-(\text{aq}) \rightleftharpoons \text{CoCl}_4^{2-}(\text{aq})$ (Blue) + $6 \text{H}_2\text{O}(\text{l})$ turns pink when placed in ice water mixture but turns blue in hot water. The reaction, as shown, is:

- A) Nonthermic B) Endothermic
 C) Exothermic D) Insufficient data

16) In which of the following aqueous solutions does the weak acid exhibit the highest percentage ionization?

- A) 0.01 M HClO ($K_a = 3.0 \times 10^{-8}$)
 B) 0.01 M HNO_2 ($K_a = 4.5 \times 10^{-4}$)
 C) 0.01 M $\text{HC}_2\text{H}_3\text{O}_2$ ($K_a = 1.8 \times 10^{-5}$)
 D) 0.01 M HF ($K_a = 6.8 \times 10^{-4}$)
 E) These will all exhibit the same percentage ionization.

17) Which of the following aqueous solutions has the highest $[\text{OH}^-]$?

- A) a solution with a pOH of 12.0
 B) a 1×10^{-3} M solution of NH_4Cl
 C) a 1×10^{-4} M solution of HNO_3
 D) a solution with a pH of 3.0
 E) pure water

18) A 0.1 M aqueous solution of _____ will have a pH of 7.0 at 25.0 °C.

18) _____

NaOCl KCl NH₄Cl Ca(OAc)₂

- A) NaOCl
- B) KCl
- C) NH₄Cl
- D) Ca(OAc)₂
- E) KCl and NH₄Cl

19) Which one of the following pairs cannot be mixed together to form a buffer solution?

19) _____

- A) NaC₂H₃O₂, HCl (C₂H₃O₂⁻ = acetate)
- B) NH₃, NH₄Cl
- C) KOH, HF
- D) H₃PO₄, KH₂PO₄
- E) RbOH, HBr

20) In which of the following aqueous solutions would you expect AgCl to have the lowest solubility?

20) _____

- A) 0.020 M BaCl₂
- B) pure water
- C) 0.020 AgNO₃
- D) 0.020 KCl
- E) 0.015 NaCl

TRUE/FALSE. On the scantron, select answer 'A' if the statement is true and 'B' if the statement is false (3 pts each).

21) Units of the rate constant of a reaction are independent of the overall reaction order.

T or F

22) In an exothermic equilibrium reaction, increasing the reaction temperature favors the formation of reactants.

T or F

23) H₂SO₃ and H₂SO₄ are considered an acid-base conjugate pair.

T or F

24) For any buffer system, the buffer capacity depends on the amount of acid and base from which the buffer is made.

T or F

25) The effect of a catalyst on a chemical reaction is to react with product, effectively removing it and shifting the equilibrium to the right.

T or F